

Table S2. Risk factor for post-UDS UTI

Variable	Post-UDS		p-value
	UTI (n=6)	No UTI (n=393)	
History of UTI within 1 month	1 (16.7)	16 (4.1)	0.231
Previous AD	0 (0)	70 (17.8)	0.596
Urinary incontinence	0 (0)	123 (31.3)	0.183
Post-void residue (>100 mL)	1 (16.7)	66 (16.8)	1.000
Pre-UDS pyuria	5 (83.3)	204 (51.9)	0.218
Pre-UDS bacteriuria	4 (66.7)	253 (64.4)	1.000
Bladder emptying method			
Voiding without catheter	2 (33.3)	122 (31.0)	1.000
Intermittent catheterization	1 (16.7)	181 (46.1)	0.226
Transurethral indwelling catheter	2 (33.3)	70 (17.8)	0.296
Suprapubic indwelling catheter	1 (16.7)	20 (5.1)	0.279
UDS findings			
Compliance (<10 mL/cmH ₂ O)	0 (0)	36 (9.2)	1.000
Maximal detrusor pressure (>40 mmHg)	1 (16.7)	116 (29.5)	0.676
Bladder capacity (mL)			
>500	6 (100)	257 (65.4)	0.790
200–500	0 (0)	125 (31.8)	0.099
<200	0 (0)	11 (2.8)	1.000

UDS, urodynamic study; UTI, urinary tract infection; AD, autonomic dysreflexia.