

Table S4. Supramaximal stimulation intensity, stratified by sex and age

	Male			Female		
	20s (n=100) ^{a)}	50s (n=100)	p-value	20s (n=100)	50s (n=98)	p-value
Tibial nerve (mA)						
Distal stimulation	54.4±12.7 (n=88)	57.4±15.9 (n=92)	0.160	53.9±16.9 (n=84)	59.3±17.5 (n=86)	0.044*
Proximal stimulation	67.0±17.0 [†] (n=86)	69.8±19.0 [†] (n=96)	0.285	70.5±19.4 [†] (n=86)	68.8±19.6 [†] (n=81)	0.573
Common peroneal nerve (mA)						
Distal stimulation	63.7±19.8 (n=85)	65.8±18.6 (n=78)	0.485	58.7±18.6 (n=79)	57.6±16.9 (n=66)	0.705
Proximal stimulation	64.1±18.5 (n=82)	66.4±17.6 (n=78)	0.421	64.1±20.4 (n=78)	59.0±16.7 (n=67)	0.103
Sural nerve (mA)	30.8±8.7 (n=93)	31.3±8.2 (n=93)	0.668	31.5±10.0 (n=91)	33.9±10.5 (n=91)	0.114
Superficial peroneal nerve (mA)	33.0±9.6 (n=93)	32.9±8.7 (n=80)	0.954	33.3±8.3 (n=90)	34.4±9.8 (n=90)	0.408

Values are presented as mean±standard deviation.

Stimulation duration for tibial nerve is 0.2 ms, and stimulation duration for other nerves is 0.1 ms.

^{a)}The total number is the number of nerve conduction studies, counting each limb separately.

*p<0.05 for Student t-test, [†]p<0.05 for Student t-test between distal and proximal stimulation.