

Supplementary Table S1. Study characteristics in SCI with TBI patients

Variable	SCI+TBI (n=74)	Concomitant TBI severity		
		Mild (n=12)	Moderate (n=40)	Severe (n=22)
Sex				
Male	58 (78.4)	7 (58.3)	32 (80.0)	19 (86.4)
Female	16 (21.6)	5 (41.7)	8 (20.0)	3 (13.6)
Age (yr)				
0-15	1 (1.3)	0 (0)	1 (2.5)	0 (0)
16-30	6 (8.1)	2 (16.7)	3 (7.5)	1 (4.5)
31-45	7 (9.5)	2 (16.7)	4 (10.0)	1 (4.5)
46-60	23 (31.1)	3 (25.0)	10 (25.0)	10 (45.5)
61-75	25 (33.8)	4 (33.3)	14 (35.0)	7 (31.8)
>75	12 (16.2)	1 (8.3)	8 (20.0)	3 (13.7)
Initial NLI				
C1-4	32 (43.2)	6 (50.0)	11 (27.5)	15 (68.2)
C5-8	14 (18.9)	2 (16.7)	11 (27.5)	1 (4.5)
T1-12	15 (20.3)	3 (25.0)	8 (20.0)	4 (18.2)
L1-S5	3 (4.1)	1 (8.3)	2 (5.0)	0 (0)
Unknown	10 (13.5)	0 (0)	8 (20.0)	2 (9.1)
Initial AIS				
A	25 (33.8)	5 (41.7)	11 (27.5)	9 (40.9)
B	3 (4.0)	0 (0.0)	1 (2.5)	2 (9.1)
C	21 (28.4)	3 (25.0)	13 (32.5)	5 (22.7)
D	15 (20.3)	4 (33.3)	7 (17.5)	4 (18.2)
Unknown	10 (13.5)	0 (0)	8 (20.0)	2 (9.1)
Traumatic etiology				
Fall	41 (55.4)	8 (66.7)	24 (60.0)	9 (40.9)
Transport	32 (43.2)	4 (33.3)	16 (40.0)	12 (54.6)
Others	1 (1.4)	0 (0)	0 (0)	1 (4.5)

Values are presented as number (%).

SCI, spinal cord injury; TBI, traumatic brain injury; NLI, neurologic level of injury; AIS, American Spinal Injury Association Impairment Scale.